

## State of Florida - Minority Health Profile - American Indian (AI)\*

Measure	Year(s)	Rate Type	Al Number	Al Rate	State Rate	Al/Total rate ratio
Socio-Demographic Characteristics						
Population <sup>1</sup>	2012		51,471	0.27%		
Population under 18 years old <sup>1</sup>	2012		10,609	20.61%		
Population 65 years and older <sup>1</sup>	2012		4,624	8.98%		
Median household income <sup>1</sup>	2012	Dollars	\$39,778		\$45,040	
Individuals below poverty level <sup>2</sup>	2008-2012 5 yr est	Percent		24.0%	17.1%	1.40:1
Civilian labor force which is unemployed <sup>1</sup>	2012	Percent		18.3%	11.5%	1.59:1
Owner-occupied housing units <sup>1</sup>	2012	Percent		51.8%	79.7%	0.65:1
Individuals 1 year and over that lived in a different house 1 year earlier <sup>1</sup>	2012	Percent		24.9%	16.1%	1.55:1
Individuals 25 years and over with no high school diploma <sup>1</sup>	2012	Percent		16.2%	13.5%	1.20:1
Access to Care						
Adults who have a personal doctor <sup>6</sup>	2012	Percent		70.9%	75.4%	0.94:1
Adults who could not see a dentist in the past year because of cost <sup>6</sup>	2010	Percent		33.9%	17.3%	1.96:1
Adults with good to excellent overall health <sup>6</sup>	2012	Percent		65.1%	80.0%	0.81:1
Adults with any type of health care insurance coverage <sup>6</sup>	2012	Percent		70.1%	78.1%	0.90:1
Maternal and Child Health						
Births <sup>3</sup>	2012	Per 1,000	303	5.9	11.0	0.53:1
Births to mothers ages 15-19 <sup>3</sup>	2012	Per 1,000	32		27.2	
Repeat births to mothers ages 15-19 <sup>3</sup>	2012	Percent	5	15.6%	17.6%	089:1
Births to mothers over 18 without high school education <sup>3</sup>	2012	Percent	38	13.4%	13.0%	1.03:1
Resident live births to mothers who smoked during pregnancy <sup>3</sup>	2012	Percent	45	14.9%	6.6%	2.25:1
Births to mothers who are overweight (BMI 25.0-29.9) at time pregnancy occurred <sup>3</sup>	2012	Percent	80	26.4%	23.8%	1.11:1
Births to mothers who are obese (BMI>=30) at time pregnancy occurred <sup>3</sup>	2012	Percent	91	30.0%	20.6%	1.46:1
Mothers who initiate breastfeeding <sup>3</sup>	2012	Percent	205	67.7%	81.4%	0.83:1
Births with 1st trimester prenatal care <sup>3</sup>	2012	Percent	228	75.2%	74.8%	1.01:1
Births with no prenatal care <sup>3</sup>	2012	Percent	5	1.7%	1.3%	1.27:1
Births with inadequate care <sup>3</sup>	2012	Percent	44	14.5%	12.6%	1.15:1
Births < 37 weeks of gestation <sup>3</sup>	2012	Percent	56	18.5%	14.3%	1.29:1
Births < 1500 grams (very low birth weight) <sup>3</sup>	2012	Percent	2	0.7%	1.6%	0.41:1
Births < 2500 grams (low birth weight) <sup>3</sup>	2012	Percent	25	8.3%	8.6%	0.96:1
Fetal deaths <sup>9</sup>	2010-2012	Per 1,000	3	3.3	7.3	0.45:1
Infant deaths (0-364 days) <sup>5</sup>	2010-2012	Per 1,000	4	4.4	6.3	0.70:1
Births with inter-pregnancy interval < 18 months <sup>3</sup>	2012	Percent	87	28.7%	20.2%	1.42:1

•

Measure	Year(s)	Rate Type	AI Number	Al Rate	State Rate	AI/Total rate ratio
Leading Causes of Death						
All cancers: Crude death rate <sup>4</sup>	2010-2012	Per 100,000	112	61.3	218.0	0.28:1
Lung cancer: Crude death rate⁴	2010-2012	Per 100,000	44	24.1	62.40	0.39:1
Colorectal cancer: Crude death rate <sup>4</sup>	2010-2012	Per 100,000	13	7.1	19.2	0.37:1
Adults 50 and over who have ever had a sigmoidoscopy or colonoscopy <sup>6</sup>	2012	Percent		47.0%	68.4%	0.69:1
Adults 50 and over who have had a blood stool test in the past year <sup>6</sup>	2012	Percent		4.2%	13.6%	0.31:1
Breast cancer: Crude death rate <sup>4</sup>	2010-2012	Per 100,000	10	5.5	15.0	0.36:1
Women 40 years of age and older who received a mammogram in the past year <sup>6</sup>	2012	Percent		57.0%	59.0%	0.97:1
Prostate cancer: Crude death rate <sup>4</sup>	2010-2012	Per 100,000	8	4.4	10.8	0.41:1
Men 50 years of age and older who received a digital rectal exam in the past year <sup>6</sup>	2010	Percent		31.2%	48.5%	0.64:1
Cervical cancer: Crude death rate <sup>4</sup>	2010-2012	Per 100,000	4	2.2	1.7	1.29:1
Women 18 years of age and older who received a PAP test in the past year <sup>6</sup>	2012	Percent		47.0%	52.9%	0.89:1
Cardiovascular Disease						
Coronary Heart Disease: Crude death rate <sup>4</sup>	2010-2012	Per 100,000	63	34.5	148.0	0.23:1
Adults who have ever had a heart attack, angina, or coronary heart disease <sup>6</sup>	2012	Percent		3.9%	8.0%	0.49:1
Stroke: Crude death rate <sup>4</sup>	2010-2012	Per 100,000	28	15.3	44.1	0.35:1
Adults who have ever had a stroke <sup>6</sup>	2012	Percent		2.6%	3.1%	0.84:1
Injuries and Injury-related Deaths						
Crude homicide death rate <sup>4</sup>	2010-2012	Per 100,000	15	8.2	6.0	1.37:1
Crude suicide death rate <sup>4</sup>	2010-2012	Per 100,000	39	21.3	14.9	1.43:1
Crude motor vehicle crash death rate⁴	2010-2012	Per 100,000	31	17.0	12.8	1.33:1
Chronic Lower Respiratory Diseases (CLRD)						
Crude death rate⁴	2010-2012	Per 100,000	37	20.2	54.6	0.37:1
Diabetes						
Crude death rate <sup>4</sup>	2010-2012	Per 100,000	26	14.2	26.6	0.53:1
Adults with diagnosed diabetes <sup>6</sup>	2012	Percent		20.7%	11.4%	1.82:1
HIV/AIDS						
Crude HIV/AIDS death rate <sup>4</sup>	2010-2012	Per 100,000	3	1.6	5.3	0.31:1
Reported AIDS Cases <sup>7</sup>	2012	Per 100,000	8	15.5	14.6	1.06:1
Reported HIV Cases <sup>7</sup>	2012	Per 100,000	14	27.2	28.3	0.96:1
Modifiable Behaviors Leading to Premature Death						
Adults who are current smokers <sup>6</sup>	2012	Percent		34.0%	17.7%	1.92:1
Adults who are obese (BMI>=30) <sup>6</sup>	2012	Percent		39.5%	25.2%	1.57:1
Adults who are overweight (BMI between 25.0 and 29.9) <sup>6</sup>	2012	Percent		32.0%	36.9%	0.87:1
Adults who engage in at least regular moderate physical activity <sup>8</sup>	2007	Percent		32.3%	34.6%	0.93:1
Adults who consume at least 5 servings of fruits and vegetables per day <sup>8</sup>	2007	Percent		28.5%	26.2%	1.09:1
Adults who have been diagnosed with high blood cholesterol <sup>6</sup>	2010	Percent		30.7%	38.6%	0.80:1

## **Notes**

\*includes solely those identifying as ONLY American Indian/Alaskan Native

Where rates are based on a sample, no number (count) is available.

## **Rate Ratios**

The rate ratios in this report compare the American Indian rate to the total Florida rate.

A ratio of 2:1 means that the American Indian rate is two times the Florida rate. A ratio of 0.5:1 means that the American Indian rate is half of the Florida rate.

## **Data Sources**

1: US Census ACS 2012 1-year estimates

http://factfinder2.census.gov/

<sup>2</sup>: US Census Bureau, ACS 2008-12 5-year estimates

http://factfinder2.census.gov/

<sup>3</sup>: Florida Birth Query System http://www.floridacharts.com/FLQuery/Birth/BirthRpt.aspx

<sup>4</sup>: Florida Death Query System http://www.floridacharts.com/FLQUERY/Death/DeathRate.aspx

<sup>5</sup>: Florida Infant Death Query System

http://www.floridacharts.com/FLQUERY/InfantMortality/InfantMortalityRpt.aspx

6: 2010 BRFSS Data

http://www.cdc.gov/brfss/

<sup>7</sup>: Florida Department of Health HIV/AIDS Bureau

8: 2007 BRFSS data

http://www.cdc.gov/brfss/

<sup>9</sup>: Florida Fetal Death Query System

http://www.floridacharts.com/FLQUERY/InfantMortality/InfantMortalityRpt.aspx